

Detection of large magneto-anisotropy of electron spin dephasing in a high-mobility two-dimensional electron system in a [001] GaAs/AlGaAs quantum well

D. Stich,¹ J. H. Jiang,² T. Korn,¹ R. Schulz,¹ D. Schuh,¹ W. Wegscheider,¹ M. W. Wu,^{2,*} and C. Schüller^{1,†}

¹*Institut für Experimentelle und Angewandte Physik,
Universität Regensburg, D-93040 Regensburg, Germany*

²*Hefei National Laboratory for Physical Sciences at Microscale and Department of Physics,
University of Science and Technology of China, Hefei, Anhui, 230026, China*

(Dated: February 6, 2008)

In time-resolved Faraday rotation experiments we have detected an inplane anisotropy of the electron spin-dephasing time (SDT) in an n -modulation-doped GaAs/Al_{0.3}Ga_{0.7}As single quantum well. The SDT was measured with magnetic fields of $B \leq 1$ T, applied in the [110] and $\bar{1}\bar{1}0$ inplane crystal directions of the GaAs quantum well. For fields along $\bar{1}\bar{1}0$, we have found an up to a factor of about 2 larger SDT than in the perpendicular direction. The observed SDTs also show strong anisotropic magnetic field dependence. Fully microscopic calculations, by numerically solving the kinetic spin Bloch equations considering the D'yakonov-Perel' and the Bir-Aronov-Pikus mechanisms, reproduce the experimental findings quantitatively. This quantitative analysis of the data allowed us to determine the relative strengths of Rashba and Dresselhaus terms in our sample. Moreover, we could predict the SDT for spins aligned in the [110] *inplane* direction to be on the order of several nanoseconds, which is up to two orders of magnitude larger than that in the perpendicular *inplane* direction.

PACS numbers: 39.30.+w 73.20.-r 85.75.-d 71.70.Ej

The exploration of electron spin relaxation and dephasing is at the heart of the research of semiconductor spintronics.^{1,2} For potential applications in quantum computation or spin transistor devices,³ the investigation and knowledge about the relevant spin dephasing channels is an important prerequisite. The most powerful techniques for the study of spin dynamics in semiconductors are all-optical techniques, like time-resolved photoluminescence or time-resolved Kerr or Faraday rotation (TRFR), where the latter are so called pump-probe techniques. In fact, TRFR experiments have helped to detect extremely long spin relaxation times in bulk GaAs structures, which were n -type doped, close to the metal-to-insulator transition.⁴ So far, however, comparatively little attention has been paid to the experimental investigation of spin dynamics in quantum wells containing free electrons (*e.g.*, in Refs. 5,6,7). In the majority of experiments, spin-aligned charge carriers are created via the absorption of circularly-polarized light. Hence, the direction of optically-oriented spins is parallel or antiparallel to the direction of light propagation. Therefore, in typical time-resolved experiments on quantum-well samples with normal incidence, the spins of the photoexcited carriers point perpendicular to the quantum-well plane, *i.e.*, in growth direction of the layered sample. In a pioneering experiment, Ohno *et al.*^{8,9} confirmed experimentally in quantum-well samples, which were grown in the [110] crystal direction, a very long spin relaxation time on the order of nanoseconds for spins, aligned parallel to the growth direction. This was predicted theoretically to be due to the absence of the D'yakonov-Perel' spin relaxation mechanism¹⁰ in this particular crystal direction. These discoveries lead to a growing interest in [110] structures, which promised to be well-suited candidates

for spintronics' devices. There are, however, essentially two important reasons, which might suggest to use standard [001]-grown heterostructures for spin-injection devices: (i) In typical real transport devices for spin injection and manipulation, as, *e.g.*, motivated by the Datta-Das spin transistor,³ layered ferromagnetic electrodes are employed as spin injectors, which exhibit inplane magnetization directions, *i.e.*, the injected spins are typically *inplane* with respect to the semiconductor structure. (ii) Due to the lower growth temperature for the growth on [110] substrates, the mobility in modulation-doped heterostructures, grown in that crystal direction, is generally lower than that in standard [001] structures.

Quite some time ago it was demonstrated theoretically that also in [001] semiconductor heterostructures of Zincblende type, spin relaxation could be greatly suppressed for spins pointing into the [110] *inplane* direction.¹³ A giant spin relaxation anisotropy was predicted for inplane spin orientation, where spin relaxation should be strong in $\bar{1}\bar{1}0$ direction and weak in the perpendicular [110] direction. The strength of the anisotropy roughly depends on the ratio between the Rashba¹¹ and Dresselhaus¹² spin-orbit coupling terms: The anisotropy should be maximal if both terms have equal strengths.¹⁴ For this case, Schliemann *et al.* proposed a spin transistor device which uses diffusive transport.¹⁵ Surprisingly, a strongly reduced spin relaxation in the [110] inplane direction in two-dimensional electron systems (2DES) in standard [001]-grown heterostructures, so far has not been detected in a time-resolved experiment. Recently, Averkiev *et al.* performed polarization-resolved cw photoluminescence measurements on a nominally undoped [001]-grown quantum well at liquid Nitrogen temperature,¹⁶ determining the depolarization of




FIG. 1: (color online) (a) Schematic picture of the time-resolved Faraday rotation experiment. Inplane magnetic fields are applied either in the $[110]$ or the $[1\bar{1}0]$ directions. (b) Sketch of the precession of optically oriented spins about a $[110]$ inplane magnetic field. (c) Same as (b) but for $[1\bar{1}0]$ magnetic field.

the photoluminescence in an applied inplane field (Hanle effect). From the experimental data the authors extracted inplane spin relaxation times, which differed by about a factor of 2 for the two perpendicular inplane directions, which showed evidence for the inplane anisotropy of the SDT. More recently, Liu *et al.* performed Kerr rotation measurement on a $[001]$ -grown two dimensional electron gas at 150 K and extracted the inplane SDTs along different directions with a factor of 1.3. In this work we report TRFR experiments on a high-mobility 2DES with inplane magnetic fields. We could detect a magneto-anisotropy of the SDT, measured via the out-of-plane component of the initial spin polarization. Calculations employing the fully microscopic kinetic spin Bloch equations,^{17,18,19,20,21} taking into account *all* relevant spin relaxation mechanisms (including both the D'yakonov-Perel' and the Bir-Aronov-Pikus mechanisms^{23,24}), reproduce the experimental data quantitatively without any parameter for the SDT. This allows us to estimate the maximum inplane SDT in our standard high-mobility quantum-well structure to be on the order of several nanoseconds, which is about 2 orders of magnitude larger than in the perpendicular inplane direction.

Our sample is a 20 nm-wide, one-sided modulation-doped GaAs-Al_{0.3}Ga_{0.7}As single quantum well. The electron density and mobility at $T = 4.2$ K are $n_e = 2.1 \times 10^{11} \text{ cm}^{-2}$ and $\mu_e = 1.6 \times 10^6 \text{ cm}^2/\text{Vs}$, respectively. For measurements in transmission geometry, the sample was glued onto a glass substrate with an optical adhesive, and the substrate and buffer layers were removed by selective etching. The sample was mounted in the ^3He insert of a superconducting split-coil magnet cryostat. All measurements were taken at a temperature of $T = 4.5$ K. For the TRFR measurements, two laser beams from a

mode-locked Ti:Sapphire laser, which is operated at 80 MHz repetition rate, were used. The laser pulses had a temporal length of about 600 fs each, resulting in a spectral width of about 3-4 meV, which allowed for a resonant excitation. The laser wavelength was tuned to excite electrons from the valence band to states slightly above the Fermi energy of the host electrons in the conduction band. Both laser beams were focused to a spot of approximately $60 \mu\text{m}$ diameter on the sample surface. The pump pulses were circularly polarized in order to create spin-oriented electrons in the conduction band, with spins aligned perpendicular to the quantum well plane, *i.e.*, in $[001]$ direction. The pump power was set to excite an initial spin polarization of the 2DES of about 6%. Such a high spin polarization ensures a long spin dephasing time to subtract the spin dephasing time from the full spin precession at low magnetic field.^{18,25} The rotation of the linear polarization of the time-delayed probe pulse, due to the Faraday effect, was measured by an optical bridge (see Fig. 1a). Due to its near-normal incidence on the sample, the polarization rotation of the probe pulse is caused by the out-of-plane component of the spin polarization of the 2DES.

Figure 1a is a schematic of the experiment, showing the orientation of the sample, relative to the laser beams and the magnetic fields. Figures 1b and 1c illustrate the precessional motion of the spin polarization in the two cases investigated in this paper: For a magnetic field applied parallel to the $[1\bar{1}0]/[110]$ direction (Fig. 1b/1c), the spins, which are created with an out-of-plane orientation parallel to the $[001]$ direction, are forced to precess within the $(1\bar{1}0)/(110)$ plane. Besides the magnetic field, the spins also precess around the momentum $\mathbf{k}$ -dependent effective magnetic field due to the Dresselhaus and Rashba spin-orbit couplings which provide an inhomogeneous broadening leading to the spin dephasing.^{14,17,18,19,20,21} Spins, precessing within the $(1\bar{1}0)/(110)$ plane, feel the inhomogeneous broadening along the $[1\bar{1}0]/[110]$ direction, which reads $(\alpha - \beta)k_x / -(\alpha + \beta)k_y$, if the y/x -axis is along $[1\bar{1}0]/[110]$.¹⁴ Here, α/β represents the Rashba/Dresselhaus coefficient. Strong anisotropy is predicted when α is comparable to β . Thus, during their precessional motion, the spins are expected to probe the relaxation times in the different inplane directions. In this work we demonstrate that the effect on their overall, averaged SDT can be observed by measuring the out-of-plane component of the spin polarization.

Figure 2 shows TRFR traces taken for various magnetic-field amplitudes applied parallel to the $[1\bar{1}0]$ direction (black traces), which show an exponentially damped oscillation of the out-of-plane component of the spin polarization. The temporal decay of the signal is determined by the SDT. A TRFR trace taken for a magnetic field of 0.9 T parallel to $[110]$ (red trace) is shown directly above for comparison. By comparing it to the corresponding trace below, the dephasing anisotropy is clearly visible. Figure 3a shows the SDTs, extracted from the experimental data by fitting an exponentially




FIG. 2: (color online) Experimental TRFR traces for different directions of the inplane magnetic field.

damped cosine function to the TRFR traces. The experimentally determined times are in excellent agreement with the calculations. A rather striking feature is the minimum in the SDT observed in Fig. 3a for a magnetic field of 0.2 T, which is clearly present in both, the experiment and the calculations. The corresponding experimental trace is shown in Fig. 3c, while Fig. 3b shows the trace at 0.2 T for the perpendicular magnetic-field direction. One can clearly see that in the latter situation the signal, and hence the out-of-plane component of the spin polarization, reflects more than half a precession cycle of the spin polarization during the measured time window, while in the former case (Fig. 3c), the spin polarization is completely relaxed within a quarter cycle. Another feature is that there is a maximum of the SDT at $B=0.5$ T for a magnetic field along $[1\bar{1}0]$ direction. The strength of the Rashba term, $\alpha = 0.65\beta$, in the calculations is tuned to generate the best fit to the experimental data, whereas the strength of the Dresselhaus term, $\beta = 1.38$ meV·Å, is determined following the work of Ref. 20.²⁶ The Rashba coefficient, $\alpha = 0.9$ meV·Å, corresponds to a built-in electric field along the z -axis of 16.8 kV/cm,²⁷ which is comparable with that of 15.0 kV/cm, which we calculated from the heterostructure of our sample self-consistently.²⁸ Using these parameters, the SDTs for spins initially aligned parallel to the magnetic field along the $[110]$ and $[1\bar{1}0]$ directions are calculated,²⁹ as Fig. 4 shows. Here, we predict that the SDT for spins aligned parallel to $[110]$ is several nanoseconds (black stars in Fig. 4), comparable to the value observed in $[110]$ -grown quantum wells.⁸

The observed phenomena can be well understood by the inhomogeneous broadening of the competing Dresselhaus and Rashba spin-orbit couplings. As the inhomoge-




FIG. 3: (color online) (a) Comparison of experimental (solid symbols) and theoretically calculated (open symbols) spin dephasing times for different inplane magnetic-field directions. α and β are the Rashba and Dresselhaus spin-orbit coefficients, respectively. (b),(c) Comparison of experimental TRFR traces at $B = 0.2$ T for the two different inplane directions of the magnetic field.




FIG. 4: (color online) Comparison of spin dephasing times, calculated with identical parameters for initial spin polarizations parallel to the $[001]$ (open symbols) and the $[110]$ and $[1\bar{1}0]$ directions (solid stars) in dependence on magnetic fields parallel to the $[110]$ and $[1\bar{1}0]$ inplane directions.

neous broadening is large/small in the $[110]$ / $[1\bar{1}0]$ direction, the spin dephasing is large/small for spin precession in the $(1\bar{1}0)$ / (110) plane, *i.e.*, along the $[1\bar{1}0]$ / $[110]$ axis. As the spin-orbit field is also strong,²⁵ a small magnetic field can not force all the spins to precess around it and there are spin precessions around the direction perpen-

dicular to the magnetic field. Therefore, increasing the magnetic field suppresses the spin precessions along the direction perpendicular to the magnetic field and hence decreases/increases the SDT when the magnetic field is along the $[110]/[\bar{1}\bar{1}0]$ direction. On the other hand, if the magnetic field is further increased, it will drive the spins to tilt toward it in favor of the energy. Due to the anisotropy of the effective magnetic field, the inhomogeneous broadening decreases/increases when spins are tilted towards the $[110]/[\bar{1}\bar{1}0]$ direction. These two competing effects lead to the minimum/maximum of the SDT observed in the experiment when the magnetic field is along $[110]/[\bar{1}\bar{1}0]$. We note that measurements with a very high accuracy of alignment of the sample with respect to magnetic field direction (less than 0.1 degree misalignment) indicate that for the magnetic field values applied in this work, there is no significant influence of a possible perpendicular field component due to slight misalignment on the SDT.

In summary, we have performed time-resolved Faraday rotation experiments on a high-mobility, $[001]$ -grown

2DES. By applying a magnetic field in two different inplane directions, we measure the inplane anisotropy of the SDT. The measurements are compared to the fully microscopic calculations and the comparison yields the strength of the Rashba spin-orbit coupling. Using these parameters, we calculate an inplane SDT of several nanoseconds for our sample. We note that even though the TRFR experiment probes the out-of-plane component of the spin polarization, it nevertheless demonstrates the inplane spin dephasing anisotropy and allows us to quantify it.

We gratefully acknowledge Jaroslav Fabian and R. T. Harley for valuable discussions. This work was supported by the Deutsche Forschungsgemeinschaft via GrK 638, grant No. Schu1171/1-3, SFB 689 and SPP1285. M.W.W. was supported by the Natural Science Foundation of China under Grant No. 10574120, the National Basic Research Program of China under Grant No. 2006CB922005, the Knowledge Innovation Project of Chinese Academy of Sciences and SRFDP. J.H.J. would like to thank J. Zhou for coding.

* Electronic address: mwwu@ustc.edu.cn.

† Electronic address: christian.schueller@physik.uni-regensburg.de.

- ¹ D. D. Awschalom, D. Loss, and N. Samarth, eds., *Semiconductor spintronics and quantum computation*, Nanoscience and technology (Springer Berlin 2002), and references therein.
- ² I. Zutic, J. Fabian, and S. Das Sarma, *Rev. Mod. Phys.* **76**, 323 (2004), and references therein.
- ³ S. Datta and B. Das, *Appl. Phys. Lett.* **56**, 665 (1990).
- ⁴ J. M. Kikkawa and D. D. Awschalom, *Phys. Rev. Lett.* **80**, 4313 (1998).
- ⁵ J. M. Kikkawa, I. P. Smorchkova, N. Samarth, and D. D. Awschalom, *Science* **277**, 1284 (1997).
- ⁶ A. Malinowski, R. S. Britton, T. Grevatt, R. T. Harley, D. A. Ritchie, and M. Y. Simmons, *Phys. Rev. B* **62**, 13034 (2000).
- ⁷ M. A. Brand, A. Malinowski, O. Z. Karimov, P. A. Marsden, R. T. Harley, A. J. Shields, D. Sanvitto, D. A. Ritchie, and M. Y. Simmons, *Phys. Rev. Lett.* **89**, 236601 (2002).
- ⁸ Y. Ohno, R. Terauchi, T. Adachi, F. Matsukura, and H. Ohno, *Phys. Rev. Lett.* **83**, 4196 (1999).
- ⁹ see also, S. Döhrmann, D. Hägele, J. Rudolph, M. Bichler, D. Schuh, and M. Oestreich, *Phys. Rev. Lett.* **93**, 147405 (2004).
- ¹⁰ M. I. D'yakonov and V. I. Perel', *Zh. Éksp. Teor. Fiz.* **60** 1954 (1971) [*Sov. Phys. JETP* **33**, 1053 (1971)].
- ¹¹ Y. A. Bychkov and E. Rashba, *Zh. Éksp. Teor. Fiz.* **39**, 66 (1984) [*Sov. Phys. JETP* **39**, 78 (1984)].
- ¹² G. Dresselhaus, *Phys. Rev.* **100**, 580 (1955).
- ¹³ N. S. Averkiev and L. E. Golub, *Phys. Rev. B* **60**, 15582 (1999); M. M. Galzov, *ibid.* **70**, 195314 (2004).
- ¹⁴ Interestingly, it is demonstrated that the longest diffusion length does not occur at exactly $\alpha = \beta$ if the cubic term of the Dresselhaus term is included by J. L. Cheng, M. W. Wu, and I. C. da Cunha Lima in *Phys. Rev. B* **75**, 205328

(2007).

- ¹⁵ J. Schliemann, J. Egues, and D. Loss, *Phys. Rev. Lett.* **90**, 146801 (2003).
- ¹⁶ N. S. Averkiev, L. E. Golub, A. S. Gurevich, V. P. Evtikhiev, V. P. Kochereshko, A. V. Platonov, A. S. Shkolnik, and Yu. P. Efimov, *Phys. Rev. B* **74**, 033305 (2006).
- ¹⁷ M. W. Wu and H. Metiu, *Phys. Rev. B* **61**, 2945 (2000).
- ¹⁸ M. Q. Weng and M. W. Wu, *Phys. Rev. B* **68**, 075312 (2003).
- ¹⁹ M. Q. Weng, M. W. Wu, and L. Jiang, *Phys. Rev. B* **69**, 245320 (2004).
- ²⁰ J. Zhou, J. L. Cheng, and M. W. Wu, *Phys. Rev. B* **75**, 045305 (2007).
- ²¹ J. L. Cheng and M. W. Wu, *J. Appl. Phys.* **99**, 083704 (2006).
- ²² B. Liu, H. Zhao, J. Wang, L. Liu, W. Wang, D. Chen, and H. Zhu, *Appl. Phys. Lett.* **90**, 112111 (2007).
- ²³ G. L. Bir, A. G. Aronov, and G. E. Pikus, *Zh. Éksp. Teor. Fiz.* **69**, 1382 (1975) [*Sov. Phys. JETP* **42**, 705 (1976)].
- ²⁴ J. Zhou and M. W. Wu, arXiv:0705.0216.
- ²⁵ D. Stich, J. Zhou, T. Korn, R. Schulz, D. Schuh, W. Wegscheider, M. W. Wu, and C. Schüller, *Phys. Rev. Lett.* **98**, 176401 (2007).
- ²⁶ Here $m_{cv} = 0.667m_0$.
- ²⁷ P. Pfeffer, *Phys. Rev. B* **59**, 15902 (1999).
- ²⁸ P. Harrison, *Quantum Wells, Wires and Dots: Theoretical and Computational Physics of Semiconductor Nanostructures* (John Wiley & Sons, 2005), Chap. 3.
- ²⁹ Unlike the SDT due to the DP mechanism calculated from the single-particle theory in the literature^{10,13} where a strong scattering limit is required, our spin kinetic Bloch equation approach is valid in both the strong and weak scattering limits. See also C. Lü, J. L. Cheng, and M. W. Wu, *Phys. Rev. B* **73**, 125314 (2006).